WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7: (11) International Publication Number: WO 00/40220 A1 A61K 9/14, 47/44, 38/13, 47/30 (43) International Publication Date: 13 July 2000 (13.07.00)

(21) International Application Number: PCT/KR99/00025 6 January 1999 (06.01.99) (22) International Filing Date:

(71) Applicant (for all designated States except US): KOREA RESEARCH INSTITUTE OF CHEMICAL TECHNOL-OGY [KR/KR]; 100, Jang-dong, Yuseong-ku, Daejeon-city 305-343 (KR).

(72) Inventors; and


- (75) Inventors/Applicants (for US only): YUK, Soon-Hong [KR/KR]; Hanbit Apt., 133-1105, Eoeun-dong, 99, Yuseong-ku, Daejeon-city 305-333 (KR). CHO, Sun-Hang [KR/KR]; Hanwool Apt., 111-404, Shinseong-dong 1-block, Yuseong-ku, Daejeon-city 305-345 (KR). LEE, Hai-Bang [KR/KR]; Hyundai Apt., 103-901, Doryong-dong, 431 Yuseong-ku, Daejeon-city 305-340 (KR).
- (74) Agent: KIM, Won-Ho; 702, Teheran Building, 825-33 Yoksam-dong, Kangnam-ku, Seoul 135-080 (KR).

(81) Designated States: JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT,

Published

With international search report.

(54) Title: METHOD OF PREPARING PHARMACEUTICAL ACTIVE INGREDIENT COMPRISING WATER-INSOLUBLE DRUG AND PHARMACEUTICAL COMPOSITION FOR ORAL ADMINISTRATION COMPRISING THE SAME



(57) Abstract

A method of preparing a pharmaceutical active ingredient comprising a water-insoluble drug is provided. In the method, a water-insoluble drug in an organic solvent is mixed with a water-soluble polymer in an aqueous solvent and spray-dried. Thereafter, the drug microparticles is mixed with oil. The method can easily prepare a pharmaceutical active ingredient including water-insoluble drug, exhibiting good bioavailability. Accordingly, it is not required for specific technology in the preparation.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Моласо	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ.	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ.	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHOD OF PREPARING PHARMACEUTICAL ACTIVE INGREDIENT COMPRISING WATER-INSOLUBLE DRUG AND PHARMACEUTICAL COMPOSITION FOR ORAL ADMINISTRATION COMPRISING THE SAME

BACKGROUND OF THE INVENTION

(a) Field of the Invention

The present invention relates to method of preparing a pharmaceutical active ingredient comprising water-insoluble drug and a pharmaceutical composition for oral administration comprising the same and, more particularly, to a method of preparing a pharmaceutical active ingredient comprising the water-insoluble improved bioavailability with simple process.

(b) Description of the Related Art

When a water-insoluble drug is orally administrated into a patient, bioavailability is low. Due to the low bioavailability, it is impossible to orally administrate the water-insoluble drug or for increasing the effective concentration of the drug in the blood, the patient frequently takes the water-insoluble drug for a long time. For example, the patient takes the water-insoluble drug three times a day for a week.

For improving bioavailability, studies to use oil and a surfactant has been attracted. Korean Patent Laid-open No. 96-21056 discloses that a drug is micro-emulsified by using oil and a surfactant. Furthermore, a drug is dissolved in an organic solvent and the mixture is micro-emulsified by using oil and a surfactant (Sandoz company, Swiss).

In addition, Korean Patent Laid-open No. 96-5136 discloses that drug in

alcohol such as ethanol or isopopanol or acetone is mixed with excess water under the condition of forming hydrosol and dried by rotary drying or freeze-drying to make preparation.

However, in the methods of Korean Patent Laid-open No. 96-21056 and Sandoz company, the surfactant used hurts gastrointestinal tract. Furthermore, the method of Korean Patent Publication No. 96-5136 has disadvantages in that manufacturing process is complicate and it is required specific manufacturing technology.

SUMMARY OF THE INVENTION

10

It is an object of the present invention to provide a method of preparing a pharmaceutical active ingredient comprising a water-insoluble drug with simple and economical process.

It is another object of the present invention to provide a pharmaceutical composition for oral administration comprising the water-insoluble drug.

These and other objects may be achieved by a method of preparing a pharmaceutical active ingredient comprising a water-insoluble drug. The method includes the steps of mixing a water-insoluble drug in an organic solvent with a water-soluble polymer in an aqueous solvent and spray-drying the mixture.

Furthermore, the present invention provides the pharmaceutical composition for oral administration including a pharmaceutical active ingredient including water-insoluble drug microparticles combined with a water-soluble polymer.

BRIEF DESCRIPTION OF THE DRAWINGS

A more complete appreciation of the invention, and many of the attendant advantages thereof, will be readily apparent as the same becomes better understood by reference to the following detailed description when considered in conjunction with the accompanying drawing, wherein:

- FIG. 1 is a graph showing water-insoluble drug excretion rate after pharmaceutical compositions including pharmaceutical active ingredients of examples 1 to 2 and comparative example 1 is oral administrated; and
- FIG. 2 is a schematic diagram showing sprayer for producing waterinsoluble drug microparticles combined with a water-soluble polymer used in the
 present invention.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a pharmaceutical active ingredient including water-insoluble drug. The pharmaceutical active ingredient exhibits good bioavailability. For that purpose, in the present invention, microparticles is formed by using a water-insoluble drug and water-soluble polymer without a surfactant which causes gastrointestinal tract damage.

The method of preparing pharmaceutical product for oral administrating
water-insoluble drug will be described in more detail.

A water-insoluble drug is dissolved in an organic solvent such as acetone to prepare a drug solution. As the water-insoluble drug, ipriflavone, biphenyl dimethyl dicarboxylate and cyclosporin may be used.

A water-soluble polymer is dissolved in an organic solvent such as

4

ethanol aqueous solution to prepare a polymer solution. As the water-soluble polymer, cellulose type compound may be used. The exemplary of cellulose type compound is ethylcellulose, polyethyleneglycol, carboxymethylcellulose, polyethyleneglycol, polyvinylpyrollidone, dextran, poloxamer and a mixture thereof.

The drug is mixed with the polymer in the weight ratio of 1:0.1 to 10, preferably 1:0.1 to 2. The mixed solution is spray-dried, thereby obtaining water-insoluble drug powder, microparticles combined with water-soluble polymer.

For obtaining microparticles, since rotary-evaporation or freeze-dry step has been used in the conventional process, the preparation is complicate, thereby requiring the specific technology. On the contrary, in the present invention, according to the performance of the spray-drying step, the preparation is simple and therefor, it is not required the specific technology.

The exemplary of a spray dryer used in the spray-drying step in the present invention is shown in Fig. 2. As shown in Fig.2, the mixed solution and air are injected into a spray-dryer body, while raising the ambient temperature by the heated air and mixed solution is changed from solution form to microparticle form by heated air.

For increasing bioavailability, the drug microparticles combined with polymer may be mixed with oil in the weight of 1:1 to 10. It is preferred that the drug microparticles mixed with oil because bioavailability increases. When oil amount is lower than 1, it is difficult to mix the drug micropartices with oil. On the contrary, oil amount exceeds 10, it is difficult to charge the mixture to a

20

5

capsule. As the oil, corn oil, peanut oil, coconut oil, caster oil, sesame oil, perilla oil, sunflower oil, walnut oil and cacao butter may be used.

The water-insoluble drug microparticles combined with water-soluble polymer may be used as an active ingredient for a pharmaceutical composition for oral administration.

The pharmaceutical composition further includes vehicle or a disintegrant, a lubricant. In addition, the pharmaceutical compositon may inlcude other pharmaceutical excipients such as stabilizer, a preservatives, an electrolyte, etc used in the pharmaceutics

The present invention is further explained in more details with reference to the following examples. The examples are not intended to limit the present invention.

10

Example 1

1 part by weight of ipriflavone was dissolved in 8 parts by weight of acetone to prepare a water-insoluble drug solution. 2 parts by weight of hydroxypropylmethylcellurose was dissolved in an aqueous ethanol solution including 1 part by weight of water and 1 part by weight of ethanol to prepare a water-soluble polymer solution. 4 parts by weight of the polymer solution was added to 9 parts by weight of the drug solution and mixed well by using a homogenizer. The mixed solution was spray-dried at 50 °C for 20 minutes by using a spray-dryer shown in Fig.2 to obtain the water-insoluble drug microparticles combined with water-soluble polymer. 3 parts by weight of the drug microparticles was mixed with 10 parts by weight of corn oil to obtain a pharmaceutical active ingredient.

6

Example 2

A pharmaceutical active ingredient was prepared by the same procedure in Example 1 except that corn oil was not used.

Comparative example 1

The commercial product of Ipriflavone was used for oral administrating water-insoluble drug.

Pharmaceutical compositions of the present invention were prepared by the conventional pharmaceutical process with these pharmaceutical active ingredients of examples 1 to 2. Furthermore, pharmaceutical compositions of comparative example 1 was prepared by disintegrating 0.27 parts by weight of Ipriflavone of comparative example 1 in water. These pharmaceutical compositions was used as powder or tablets. These tablets or powder were oral administered into white rats in the amount of 50 mg of drug per kg of rat's weight with conventional method. The amounts of ipriflavone in the tablets or powder which did not absorbed in the rat's body and excreted to the out of the body were determined. The result shows in Fig. 1.

The result of comparative example 1 show that the most of ingredient ipriflavone did not absorbed in the rat's body and excreted to the out of the body were shown in Fig. 1. As shown in Fig. 1, when the pharmaceutical active ingredient including the drug microparticles and water of the Example 2 was oral administered, absorption in the body increased two times more than the Comparative example 1. Furthermore, when the pharmaceutical active ingredient including oil of example 1, absorbing in the body is seven times more than the comparative example 1.

7

As described above, when the drug microparticles obtained by the present invention is oral administered with oil, the increased bioavailability is exhibited. Accordingly, the drug of the present invention can be oral administered. Furthermore, it is not required for the drug to administered into a patient three or four times a day rather than the conventional water-insoluble drug.

While the present invention has been described in detail with reference to the preferred embodiments, those skilled in the art will appreciate that various modifications and substitutions can be made thereto without departing from the spirit and scope of the present invention as set forth in the appended claims.

WHAT IS CLAIMED IS:

15

20

- 1. A method of preparing a pharmaceutical active ingredient composition including water-insoluble drug for oral administration comprising the steps of:
- mixing water-insoluble drug in an organic solvent with water-soluble polymer in an aqueous solvent;

spray-drying the mixture.


- 2. The method of claim 1 wherein the polymer is selected from the group consisting of ethylcellulose, polyvinylpyrrolidone, polyethyleneglycol,

 carboxymethylcellulose, hydroxypropylmethylcellurose, dextran and poloxamer.
 - 3. The method of claim 1 wherein the drug is selected from the group consisting of ipriflavone, biphenyldimethyldicarboxylate and cyclosporin.
 - 4. The method of claim 1 wherein the weight ratio of the drug and polymer is 1 : 0.1 to 10.
 - 5. The method of claim 1 further comprising the step of mixing the microparticles with oil.
 - 6. The method of claim 1 wherein the oil is selected from the group consisting of corn oil, peanut oil, coconut oil, castor oil, sesame oil, perilla oil, sunflower oil, walnut oil and cacao butter.
 - 7. The method of claim 5 wherein the weight ratio of the microparticles and oil is 1 : 1 to 10.
 - 8. A pharmaceutical composition for oral administration, comprising a pharmaceutical active ingredient water-insoluble drug microparticles combined with water-soluble polymer.

9

- 9. The pharmaceutical composition of claim 8 wherein the drug is selected from the group consisting of ipriflavone, biphenyldimethyldicarboxylate and cyclosporin.
- 10. The pharmaceutical composition of claim 8 wherein the polymer is selected from the group consisiting of ethylcellulose, polyvinylpyrrolidone, polyethyleneglycol, carboxymethylcellulose, hydroxypropylmethylcellurose, dextran and poloxomer.

FIG.1



Outlet Outlet

Spray-dryer body

Heated air

1/1

INTERNATIONAL SEARCH REPORT

International application No. PCT/KR 99/00025

A. CLASSIFICATION OF SUBJECT MATTER

IPC⁷: A 61 K 9/14, A 61 K 47/44, A 61 K 38/13, A 61 K 47/30

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: A 61 K, C 08 L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, TXTE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Х	EP 0 670 166 A2 (ELI LILLY COMPANY) 06 September 1995 (06.09.95), abstract; page 5, lines 2-10; claims 1, 2, 5, 8.	1-10
Х	WO 95/32726 A (YUHAN CORPORATION) 07 December 1995 (07.12.95), abstract; page 4, lines 15-25; claims.	1-4, 8-10
X	US 5 817 343 A (BURKE) 06 October 1998 (06.10.98), abstract; column 6, lines 63.67; column 7, lines 1-15, 40-45; claims 1, 31, 33.	1,2,8,10
X	EP 0 012 496 A1 (BEECHAM GROUP LIMITED) 25 June 1980 (26.06.80), abstract; page 3, lines 1-16; page 4, lines 18-21; page 5, lines 1-12.	1,2,4,8

Further documents are listed in the continuation of Box C.	See patent family annex.
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
Date of the actual completion of the international search	Date of mailing of the international search report
14 October 1999 (14.10.99)	08 November 1999 (08.11.99)
Name and mailing adress of the ISA/AT	Authorized officer
Austrian Patent Office	Krenn
Kohlmarkt 8-10; A-1014 Vienna	130im
Facsimile No. 1/53424/200	Telephone No. 1/53424/435

Telephone No. 1/53424/435

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR 99/00025

Doc ange	le Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche		Datum der Veröffentlichung Publication state Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets		Datum der Veröffentlichung Publication date Date de publication
EF	A2	57 01 <i>5</i> 5	06-09-1995	ARANAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA	13544/95 950060784 21386088 950706189 450706182 950752745 80450454 9501344 9501469 9501469 9501469 9507936	07-09-1995 24-10-1995 024-09-19995 10-10-19995 10-10-19995 10-04-19995 28-05-19995 28-05-19995 13-01-19995 13-019-19995 04-08-1995 09-10-1995
WO.	A1	9532726		keine -	none - rie	3n
นธ	A	5817343	06-10-1998	AU A1 CA AA EP A1 WO A1	27517/97 2253667 914095 9742940	05-12-1997 20-11-1997 12-05-1999 20-11-1997
EP	A1	1.2496	25-06-1980	AU A1 AU B2 DE CO EP B1 JP A2 NX A	52284/79 531634 2965917 12496 55079387 191982 4316886	12-06-1980 01-09-1983 25-08-1983 20-07-1983 14-06-1980 23-11-1982 23-02-1982

Form PCT/ISA/210 (patent family annex) (July 1998)